

2. Assay For Synthetic Activity

The ability of the modified enzyme or proteolytic fragments is assayed by adding the modified enzyme to an assay system in which a primer is annealed to a template and DNA synthesis is catalyzed by the added enzyme. Many standard laboratory techniques employ such an assay. For example, nick translation and enzymatic sequencing involve extension of a primer along a DNA template by a polymerase molecule.

In a preferred assay for determining the synthetic activity of a modified enzyme an oligonucleotide primer is annealed to a single-stranded DNA template, *e.g.*, bacteriophage M13 DNA, and the primer/template duplex is incubated in the presence of the modified polymerase in question, deoxynucleoside triphosphates (dNTPs) and the buffer and salts known to be appropriate for the unmodified or native enzyme. Detection of either primer extension (by denaturing gel electrophoresis) or dNTP incorporation (by acid precipitation or chromatography) is indicative of an active polymerase. A label, either isotopic or non-isotopic, is preferably included on either the primer or as a dNTP to facilitate detection of polymerization products. Synthetic activity is quantified as the amount of free nucleotide incorporated into the growing DNA chain and is expressed as amount incorporated per unit of time under specific reaction conditions.

Representative results of an assay for synthetic activity is shown in Figure 18. The synthetic activity of the mutant DNAP_{Taq} clones 4B-F was tested as follows: A master mixture of the following buffer was made: 1.2X PCR buffer (1X PCR buffer contains 50 mM KCl, 1.5 mM MgCl₂, 10 mM Tris-Cl, pH 8.5 and 0.05% each Tween 20 and Nonidet P40), 50 μM each of dGTP, dATP and dTTP, 5 μM dCTP and 0.125 μM α-³²P-dCTP at 600 Ci/mmol. Before adjusting this mixture to its final volume, it was divided into two equal aliquots. One received distilled water up to a volume of 50 μl to give the concentrations above. The other received 5 μg of single-stranded M13mp18 DNA (approximately 2.5 pmol or 0.05 μM final concentration) and 250 pmol of M13 sequencing primer (5 μM final concentration) and distilled water to a final volume of 50 μl. Each cocktail was warmed to 75°C for 5 minutes and then

cooled to room temperature. This allowed the primers to anneal to the DNA in the DNA-containing mixtures.

For each assay, 4 μ l of the cocktail with the DNA was combined with 1 μ l of the mutant polymerase, prepared as described, or 1 unit of DNAP*Taq* (Perkin Elmer) in 1 μ l of dH₂O. A "no DNA" control was done in the presence of the DNAP*Taq* (Figure 18, lane 1), and a "no enzyme" control was done using water in place of the enzyme (lane 2). Each reaction was mixed, then incubated at room temperature (approx. 22°C) for 5 minutes, then at 55°C for 2 minutes, then at 72°C for 2 minutes. This step incubation was done to detect polymerization in any mutants that might have optimal temperatures lower than 72°C. After the final incubation, the tubes were spun briefly to collect any condensation and were placed on ice. One μ l of each reaction was spotted at an origin 1.5 cm from the bottom edge of a polyethyleneimine (PEI) cellulose thin layer chromatography plate and allowed to dry. The chromatography plate was run in 0.75 M NaH₂PO₄, pH 3.5, until the buffer front had run approximately 9 cm from the origin. The plate was dried, wrapped in plastic wrap, marked with luminescent ink, and exposed to X-ray film. Incorporation was detected as counts that stuck where originally spotted, while the unincorporated nucleotides were carried by the salt solution from the origin.

Comparison of the locations of the counts with the two control lanes confirmed the lack of polymerization activity in the mutant preparations. Among the modified DNAP*Taq* clones, only clone 4B retains any residual synthetic activity as shown in Figure 18.

EXAMPLE 3

5' Nucleases Derived From Thermostable DNA

Polymerases Can Cleave Short Hairpin Structures With Specificity

The ability of the 5' nucleases to cleave hairpin structures to generate a cleaved hairpin structure suitable as a detection molecule was examined. The structure and sequence of the hairpin test molecule is shown in Figure 19A (SEQ ID NO:15). The

oligonucleotide (labeled "primer" in Figure 19A, SEQ ID NO:22) is shown annealed to its complementary sequence on the 3' arm of the hairpin test molecule. The hairpin test molecule was single-end labeled with ³²P using a labeled T7 promoter primer in a polymerase chain reaction. The label is present on the 5' arm of the hairpin test molecule and is represented by the star in Figure 19A.

The cleavage reaction was performed by adding 10 fmoles of heat-denatured, end-labeled hairpin test molecule, 0.2uM of the primer oligonucleotide (complementary to the 3' arm of the hairpin), 50 μM of each dNTP and 0.5 units of DNAP*Taq* (Perkin Elmer) or 0.5 μl of extract containing a 5' nuclease (prepared as described above) in a total volume of 10 μl in a buffer containing 10 mM Tris-Cl, pH 8.5, 50 mM KCl and 1.5 mM MgCl₂. Reactions shown in lanes 3, 5 and 7 were run in the absence of dNTPs.

Reactions were incubated at 55° C for 4 minutes. Reactions were stopped at 55° C by the addition of 8 μl of 95% formamide with 20 mM EDTA and 0.05% marker dyes per 10 μl reaction volume. Samples were not heated before loading onto denaturing polyacrylamide gels (10% polyacrylamide, 19:1 crosslinking, 7 M urea, 89 mM Tris-borate, pH 8.3, 2.8 mM EDTA). The samples were not heated to allow for the resolution of single-stranded and re-duplexed uncleaved hairpin molecules.

Figure 19B shows that altered polymerases lacking any detectable synthetic activity cleave a hairpin structure when an oligonucleotide is annealed to the single-stranded 3' arm of the hairpin to yield a single species of cleaved product (Figure 19B, lanes 3 and 4). 5' nucleases, such as clone 4D, shown in lanes 3 and 4, produce a single cleaved product even in the presence of dNTPs. 5' nucleases which retain a residual amount of synthetic activity (less than 1% of wild type activity) produce multiple cleavage products as the polymerase can extend the oligonucleotide annealed to the 3' arm of the hairpin thereby moving the site of cleavage (clone 4B, lanes 5 and 6). Native DNAP*Taq* produces even more species of cleavage products than do mutant polymerases retaining residual synthetic activity and additionally converts the hairpin structure to a double-stranded form in the presence of dNTPs due to the high level of synthetic activity in the native polymerase (Figure 19B, lane 8).